



"Tropical" diversity of *Penicillium* spp. in water-related polar environments

Sonjaka, Silva; Frisvad, Jens Christian; Gunde-Cimermana, Nina

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):

Sonjaka, S., Frisvad, J. C., & Gunde-Cimermana, N. (2011). *"Tropical" diversity of Penicillium spp. in water-related polar environments*. Abstract from Polar and Alpine Microbiology, Ljubljana, Slovenia, Ljubljana, Slovenia. <http://www.polaralpinemicrobiology.com/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

“Tropical” diversity of *Penicillium* spp. in water-related polar environments

Silva Sonjak^a, Jens Christian Frisvad^b, Nina Gunde-Cimerman^{a,c}

^a*Department of Biology, Biotechnical Faculty, University of Ljubljana, Večna pot 111, SI-1000 Ljubljana, Slovenia;*

^b*Department of Systems Biology, Technical University of Denmark, Søltofts Plads, Building 221, DK-2800 Kgs. Lyngby, Denmark;*

^c*Centre of Excellence for Integrated Approaches in Chemistry and Biology of Proteins (CIPKeBiP), Jamova 39, SI-1000 Ljubljana, Slovenia.*

E-mail: silva.sonjak@bf.uni-lj.si

Keywords: filamentous fungi, distribution, *Penicillium*, sea water, Svalbard

Penicillium species are not only amongst the most frequently isolated filamentous fungi from polar and alpine soil, but also from water environments. An almost ‘tropical’ diversity and quantity of *Penicillium* spp. was isolated in the study of the water-related coastal Arctic environments of the Svalbard archipelago in years 2001, 2003 and 2008. All together, at least 35 different *Penicillium* species were recorded. Most of them (78%) originated from different glacial ice samples. The highest numbers of isolated strains and the highest species diversity were obtained from (sub) glacial ice of different glaciers. A considerably lower diversity of *Penicillium* species was detected as well in glacial melt water, bordering sea ice and sea water, and puddles on snow in the tidal zone. Most species released from the glaciers into the surrounding area were detected in considerably lower numbers, however certain species, like *P. nordicum*, were isolated from sea water in high numbers. In the distribution of conidial filamentous fungi air is most often considered as the main dissemination factor, however recent evidence based on fungi from extreme environments, including *Penicillium* spp. indicates that water environments, such as glacial ice and sea-water might contribute much more than previously assumed to the distribution of certain pan-global species, particularly those that are involved in contamination of our house-holds and our food.